FORM PTO-1390 U.S. DEPARTMENT OF COM	MERCE PATENT AND TRADEMARK OFFICE	ATTORNEY 'S DOCKET NUMBER
(REV. 1-98)		MSKP039
TRANSMITTAL LETTER	TO THE UNITED STATES	U.S. APPLICATION NO (If known, see 37 CFR 1 5
CONCERNING A FILIN	ED OFFICE (DO/EO/US) IG UNDER 35 U.S.C. 371	09/674975
INTERNATIONAL APPLICATION NO PCT/US99/10065	INTERNATIONAL FILING DATE 07 May 1999	PRIORITY DATE CLAIMED 08 May 1998
TITLE OF INVENTION COMPOSITIONS AND METHODS FOR	ACTIVE VACCINATION	EL556132244U
APPLICANT(S) FOR DO/EO/US		FF33873EE440
Agus, et al. Applicant herewith submits to the United St	ates Designated/Elected Office (DO/EO/US	b) the following items and other information:
	s concerning a filing under 35 US C 371	
	NT submission of items concerning a filing	; under 35 US.C 371
This express request to begin nation examination until the expiration of the	al examination procedures (35 US.C 371(f) the applicable time limit set in 35 U S C371) at any time rather than delay (b) and PCT Articles 22 and 39(1)
4. A proper Demand for International Pr	reliminary Examination was made by the 19t	h month from the earliest claimed priority date
	ion as filed (35 U S.C.371(c)(2))	(i al Parana)
= 1 1	quired only if not transmitted by the International Bureau	itional Bureau).
b. Inas been transmitted by in c. In is not required, as the appl	ication was filed in the United States Recei	ving Office (RO/US)
A translation of the International Ap	oplication into English (35 U.S C 371(c)(2)	Į.
	ternational Aplication under PCT Article 19	
a. are transmitted herewith (required only if not transmitted by the Intern	national Bureau).
b. have been transmitted by the have not been made; however	the International Bureau ever, the time limit for making such amendi	ments has NOT expired
d. A have not been made; howe		none has a co-part
8. A translation of the amendments to	the claims under PCT Article 19 (35 U S C	371 (c)(3)).
9. An oath or declaration of the invente		
i 7	nternational Preliminary Examination Repo	ort under PCT Article 36
Items 11. to 16 below concern docume	ent(s) or information included:	
11. An Information Disclosure Stateme	nt under 37 CFR 1.97 and 1.98.	
12. An assignment document for record	ling A separate cover sheet in compliance	with 37 CFR 3 28 and 3.31 is included
13. A FIRST preliminary amendment		
☐ A SECOND or SUBSEQUENT pre	eliminary amendment.	
14. A substitute specification.		
15. A change of power of attorney and/	or address letter	
16. Other items or information:		Express Mail Cert. Exp. Mail No.: EL 556133344US Date Mailed: OT November 2000 Name: SUSAN HOWARD Sign:
		aign: Swen Howark

528 Rec'd PCT/PTO **0** 7 NOV 2000

U.S. APPLICATION NO (1fd.nov	vi. sec 37 CFR 15) 7 R PCT/U	TERNATIONAL APPLICATION NO S99/10065			ATTORNEY'S DOCKE MSKP039	T NUMBER
					LCULATIONS P	TO USE ONLY
BASIC NATIONAL FEE (37 CFR 1 492 (a) (1) - (5)):						
Neither internation	al preliminary examinati	ion fee (37 CFR 1.482)				
nor international se and International Se	arch fee (37 ČFR 1.445(a earch Report not prepare	a)(2)) paid to USPTO d by the EPO or JPO	\$1070 00			
International prelin USPTO but Interna	ninary examination fee (I ntional Search Report pre	37 CFR 1 482) not paid to epared by the EPO or JPC	o) \$930 00			
International prelimbut international se	ninary examination fee (2 arch fee (37 CFR 1 445(a	37 CFR 1 482) not paid to a)(2)) paid to USPTO	O USPTO \$790 00			
but all claims did r	not satisfy provisions of I	37 CFR 1.482) paid to US PCT Article 33(1)-(4)	(\$720.00			
International prelin	ninary examination fee (I fied provisions of PCT A	37 CFR 1.482) paid to U	SPTO \$98.00			
		BASIC FEE AMO		\$	110000	
			□ 20 □ 30	-	710,00	
months from the ear	0 for furnishing the oath liest claimed priority dat	e (37 CFR 1 492(e))		\$		
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$		
_Total claims	∂ O -20 =	0	x \$22.00	\$		
findependent claims	4 -3 =		x \$82.00	\$	89.00	
The Control of the Co	DENT CLAIM(S) (if app		+ \$270.00	\$ \$	700	
5106		OF ABOVE CALCU applicable. A Small Ent			792,00	
must also be filed (1	Note 37 CFR 1.9, 1.27, 1.	(28) Applicant clair See 37 CFR	ms small entity status +	\$ -	396.00	
<u>.</u>			JBTOTAL =	\$	396,00	
Processing fee of \$1	30.00 for furnishing the	English translation later	than 20 30	\$		
months from the ear	liest claimed priority dat			<u> </u>		
		TOTAL NATIO		\$	396.00	
Fee for recording the accompanied by an	e enclosed assignment (3 appropriate cover sheet (7 CFR 1.21(h)) The ass 37 CFR 3 28, 3.31) \$40	signment must be 1.00 per property +	\$		
		TOTAL FEES E	NCLOSED =	\$	396.00	
101111111111111111111111111111111111111				Am	ount to be refunded:	\$
					charged:	\$
b. 🗂 Please cha	rge my Deposit Account e copy of this sheet is en	No i	ne above fees is enclo		to cover the	above fees.
c. ☑ The Comm	nissioner is hereby autho	rized to charge any addit o. <u>15-061</u> 0 A dupli	ional fees which may cate copy of this shee	be re	equired, or creditanclosed.	any
NOTE: Where at 1.137 (a) or (b)) m	n appropriate time limi oust be filed and granted	t under 37 CFR 1.494 or d to restore the applicat	r 1.495 has not been ion to pending status	met,	a p e ition to reviv	re (37 CFR
SEND ALL CORRESP	ONDENCE TO		Mar	us	u S X	as
			SIGNATI	URE		
			Marina	T. L	arson	
02	1121		NAME			
PATENT TRA	DEMARK OFFICE		32,038	3		
			REGISTE	RATIO	N NUMBER	
					•	

25

5

PCT/US99/10065

COMPOSITIONS AND METHODS FOR ACTIVE VACCINATION

BACKGROUND OF THE INVENTION

This application relates to an active vaccine approach to the treatment of cancer and other diseases. The approach is applicable to a number of cancers and diseases, although a preferred embodiment provides an active vaccine for treatment of B cell Non-Hodgkin's Lymphoma (NHL).

NHL is characterized by a clonal proliferation of malignant B cells. The treatment of NHL across a broad spectrum of patients remains a challenge, although numerous therapeutic approaches have been proposed and tried.

The most common therapeutic approach being used today is chemotherapy. While chemotherapy is effective for some period of time in most patients, a significant percentage of patients are not cured and experience a relapse.

Treatments have been proposed based on anti-idiotype therapy. In anti-idiotype therapy, a cell surface molecule which is expressed by malignant cells but not by normal cells is used to create patient-specific antibodies which are then administered to the patient. See, Miller, et al., *New Engl. J. Med.* 306: 517-522 (1982). Autologous patient-derived idiotype proteins have also been conjugated with keyhole limpet hemocyanin to produce a vaccine which has demonstrated efficacy and can elicit B and T cell immune responses. Kwak et al., *New Engl. J. Med.* 327: 1209-1215 (1992). Hybridoma-derived idiotype was co-cultured with patient-derived dendritic cells which acted as antigen presenters upon re-infusion into the patient and showed clinical efficacy. Hsu et al., *Nature Medicine* 2: 52-58 (1996). Idiotypic vaccines made in lipid-based carriers are disclosed in International Patent Publication WO98/14170.

Treatments have also been proposed using antibodies directed to CD20, a transmembrane protein that is expressed by both normal and malignant B-cells during parts of the B cell development cycle. Using single-dose infusions with anti-CD20 monoclonal antibodies, partial or minor tumor regressions were observed in 6 of 15 patients in a Phase I clinical study. Maloney et al., *Blood* 84: 2457-2466 (1994). In Phase II studies, 17 of 37 patients showed complete or partial remissions. In December 1997, the FDA approved the

first antibody-based therapy for NHL. Rituximab (Ritvaxan, IDEC/Genentech) is a chimeric human/murine antibody approved for the treatment of patients with relapsed or refractory low-grade or follicular CD20⁺ B cell NHL. Maloney et al., *Blood* 90: 2188-2195 (1997).

Combinations of chemotherapy and anti-CD20 therapy have been reported as having better therapeutic efficacy, with 11 of 11 patients showing complete or partial remission. Czuczman et al., Abstract 53, Ann. Oncol. 7, Supp. 1: 56 (1996).

While therapeutic regimens using anti-CD20 concepts are potentially effective, all of these therapies have the drawback of being passive therapies, i.e., they do not directly involve the immune system of the patient. Thus, these therapies may require the continued administration of the therapeutic agent for efficacy and do not provide any long-term protection against recurrence. In addition, the passive therapy is monoclonal in nature, therefore escape is possible. It would therefore be desirable to have an active therapy, that is a therapeutic agent which when administered to the patient stimulates an immune response against CD20 found in B-cells.

It is an object of the present invention to provide such a therapy. It is a further object of the invention to provide an active polyclonal therapy that is difficult to evade.

SUMMARY OF THE INVENTION

20

In accordance with the present invention, NHL is treated, not by administration of an anti-CD20 monoclonal antibody, but by the administration of CD20 itself, or an immunogenic fragment of the extracellular portion thereof, coupled to or administered with an antigenic carrier moiety such as keyhole limpet hemocyanin (KLH). This results in the stimulation of the production of polyclonal antibodies against CD20 (or an immunogenic fragment thereof) which has the affect of reducing the number of B-cells, including malignant B-cells. Thus, the invention provides an active vaccine. The same approach can be used for therapeutics for other diseases and conditions in which target cells possess a transmembrane protein, and is particularly applicable to those diseases and conditions for which administration of antibodies to transmembrane proteins or peptides (i.e., passive therapy) have been shown to provide therapeutic benefits, and especially in the

25

30

5

situations where the target is also capable of transducing or receiving a signal important for cell growth or function. This would include, for example, Her2/neu, VEGF receptor, epidermal growth factor receptor, the CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-glycoprotein, also known as the multidrug-resistance protein.

BRIEF DESCRIPTION OF THE FIGURES

Figs. 1A and B show ELISA results for formation of antibodies to human and mouse CD20 in vaccinated mice;

Figs. 2A and B shows results for binding of control B1 antibodies or antibodies in plasma from a mouse treated with human CD20-KLH conjugate with Raji B NHL cells;

Fig. 3 shows CP19⁺B cell levels in mice treated with human or mouse CD20-KLH conjugate;

Fig. 4 shows the domain structure of human Her2;

Fig. 5 shows the domain structure of human EGFR;

Figs. 6A-D shows the cross-reactivity of antibodies generated in response to human or mouse CD20 fragments;

Figs. 7A-D show the importance of carrier protein and adjuvant in generating an immune response;

Figs. 8A-D shows the immune response generated using different adjuvants; and

Figs. 9A-I shows CP19⁺B cell levels in mice treated with human or mouse CD20-KLH conjugate.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides an active vaccine therapy which can be used in the treatment of a variety of cancers and related conditions in which it is desirable to bring about the death of a target group of cells. Conventionally, immunotherapies targeting cells have sought to obtain a cellular immune response (T-cells that recognize the target cells), since a humoral immune response (antibodies that recognize the target cells) alone is not

5

deemed sufficient to achieve the desired result of cell death. The present invention departs from this conventional wisdom, and effectively utilizes a humoral immune response against the target cells to provide therapeutic benefit. The targets for therapy include cell surface proteins that when bound by a ligand signal to the cell. The vaccine induced antibody response will mimic ligand binding and cause similar signaling events which can imitate the process of programmed cell death (apoptosis) or halt the cell from growing or change the cancer cell's sensitivity to chemotherapy.

By way of example, the invention is suitably employed in the treatment of NHL and other B cell diseases such as chronic lymphocytic leukemia, auto-immune disorders and B-cell regulatory disorders. In accordance with this embodiment of the invention, a peptide antigen is prepared which contains at least an immunogenic portion of the extracellular domain of CD20 coupled to or administered with an antigenic carrier protein. The CD20 component of the peptide antigen may be syngeneic or it may be xenogeneic. Thus, for example, human patients may be treated with a peptide vaccine containing a human or a mouse CD20-fragment. There is evidence that strong immune responses can be elicited against xenogeneic proteins. Naftzger et al., Proc. Natl. Acad. Sci. (USA) 93: 14809-14814 (1996); International Patent Application PCT/US97/22669, filed December 10 1997, incorporated herein by reference. A suitable fragment is the 44 amino acid peptide spanning amino acids 136 to 179 of the sequence of mouse or human CD20. (Seq. ID Nos. 1 and 2) Other immunogenic fragments derived from the extracellular domain of CD20, or the entire CD20 molecule may also be used. Seq. ID. Nos. 3 and 4 shows the nucleic acid and amino acid sequences, respectively, of exon VI (the extracellular domain) of human CD20 as reported by Tedder et al., J. Immunol. 142: 2560-2568 (1989).

As used in the specification and claims hereof, an "immunogenic fragment" is a molecule which includes at least a portion of the extracellular domain of a transmembrane protein to direct and immunological response to that transmembrane protein when the immunogenic fragment is coupled to or administered with an antigenic carrier protein effective to break tolerance and administered with an adjuvant. It is not required that the immunogenic fragment alone be effective to stimulate an immune response, although such stimulation would not take a given fragment outside the scope of the present invention.

25

5

A preferred antigenic carrier protein is keyhole limpet hemocyanin which can be coupled to peptides using techniques described in Pierce Catalog Protocol. Other antigenic carrier proteins which can be used to break tolerance might be used in the invention include immunoglobulins, tuberculin, tetanus toxin and others well known in the art.

The peptide antigen containing the CD20 component and the antigenic carrier protein is formulated with a pharmaceutically acceptable adjuvant in a liquid carrier and administered to a patient suffering from NHL or another B cell disease. The composition will generally be administered by injection, for example, intramuscular, subcutaneous or intradermal injection, but might also be administered by way of a DNA vaccine (See US Patent No. 5,580,859, incorporated herein by reference) or a viral vaccine, or after mixing with antigen presenting cells (APC's) such as dendritic cells, *ex vivo*. Alternatively, the antigen may be administered without adjuvant by injection into a host prepared by prior or simultaneous injection of an immune adjuvant. Specific amounts to be administered to a patient can be determined by monitoring the titer of anti-CD20 antibodies developed by the patient, or by an average group of patients using well-known technology.

When a peptide of the extracellular domain of human or mouse CD20 is coupled to KLH and administered with an adjuvant to mice, antibodies which react with CD20 are found in plasma. (Figs. 1A and B) These antibodies bind to Raji cells, a human lymphoma cell line, indicating the ability to bind to a cell expressing CD20. (Figs. 2A and B). Moreover, the number of CD19⁺ B cells present in mice injected with either of the two CD20-KLH conjugates declines substantially (~30% decrease relative to controls). (Figs. 3 and 9). The assay used to quantitate B cell depletion detects CD19 which is also expressed on immature B cells that are CD20⁻. Thus, the 30% depletion actually underestimates the efficacy of the vaccine against CD20⁺ B cells.

Antibodies generated in mice after vaccination with human or mouse-derived CD20 fragments are specific for the peptides used, yet are capable of inducing immunity to the corresponding peptide from other species (Figs. 6A-D). Studies showed that in most instances the peptide, carrier protein and adjuvant are all needed for optimal response, although some responses were detected using less than all of the components. (Figs. 7A-D).

25

30

5

Several different adjuvants were also tested, and QS21 was found to be the most effective of those tested. (Fig. 8A-D).

While not intending to be bound by any particular mechanism, it is believed that the vaccines of the present invention are effective via at least two pathways. First, the generation of a humoral immune response to CD20 is effective to some extent to reduce the numbers of B cells bearing CD20 antigen in a manner consistent with normal immunological response to a target antigen. In addition, however, because CD20 has a signaling function, the binding of antibody to the CD20 moiety activates this signaling function to trigger apoptotic cell death. Such stimulation of apoptosis has been demonstrated to occur *in vitro* following passive treatments with a chimeric anti-CD20 antibody. Maloney et al., *Blood* 88 (Supp. 1): 637a (1996).

It is also possible that T cell mediated effector mechanisms are involved in the immune response. As evidence of this, we illustrate in Table 1 the mouse and human peptide sequences capable of binding to the corresponding mouse and human histocompatability antigens. This information was derived from a search of the NIH Bioinformatics and Molecular Analysis Section HLA Binding Predictions database using the mouse and human CD20 amino acid sequences. (Parker et al., *J. Immunol.* 152: 163 (1994)).

While the method of the invention is illustrated here using CD20 or CD20-derived peptides as the antigen to target B cells, the invention is not limited to this embodiment. Rather, the inventions encompasses the use of vaccine compositions comprising an immunogenic portions of the extracellular domain of transmembrane protein or peptide, particularly a transmembrane protein or peptide having signaling function, coupled to or administered with an antigenic protein and/or adjuvant to break tolerance.

A non-limiting example of another transmembrane protein which can be used in whole or in part in the method of the invention is Her-2/neu. The Her-2/neu oncogene is a receptor-like tyrosine kinase that is expressed on the cell surface of a significant portion of solid tumors. It has been shown that patients with early stage breast cancer have a high titer of antibodies to Her-2/neu. Disis et al., *J. Clin. Oncol.* 15: 3363-3367 (19967). The amino acid sequence and domain structure of human Her-2/neu are shown in Seq. OD. No. 5 and Fig. 4, and isolation and expression of the extracellular domain has been disclosed.

25

20

International Patent Publication No. WO 90/14357, which is incorporated herein by reference. There is clinical data showing efficacy of monoclonal antibodies against Her-2-neu in the treatment of patients with Her-2/neu⁺ tumors, and potential synergism with chemotherapy. Thus, in accordance with the present invention, a vaccine composition comprising at least an immunogenic portion of the extracellular domain of Her-2-neu (amino acids 22 to 652) coupled to or administered with an antigenic protein or peptide such a KLH can be used as a vaccine to provide the same therapeutic benefits using an active as opposed to a passive approach.

A further non-limiting example of a transmembrane protein which can be used in whole or in part in the method of the invention is epidermal growth factor receptor (EGFR). The amino acid sequence and domain structure of human EGFR are shown in Seq. ID. No. 6 and Fig. 5. There is significant data showing that antibodies to EGFR can have anti-tumor activity in breast and prostate cancer, as well as several head and neck tumors. Prewett et al., *J. Immunother. Emphasis Tumor Humoral* 19: 419-27 (1996). The mechanism by which antibody therapy against EGFR may be efficacious can be through the ability to down-regulate vascular endothelial growth factor production by tumor cells and thereby decrease angiogenesis. Petit et al., *Am. J. Pathol.* 151: 1523-30 (1997). In accordance with the present invention, a vaccine composition comprising at least an immunogenic portion of the extracellular domain of EGFR (amino acids 25 to 645) coupled to or administered with an antigenic protein or peptide such a KLH can be used as a vaccine to provide the same therapeutic benefits using an active as opposed to a passive approach. Preferred immunogenic peptides would be selected from regions not deleted in the various types of truncated EGFR mutants associated with some cancers.

A further non-limiting example of a transmembrane protein which can be used in whole or in part in the method of the invention is VEGF receptor. There are significant data showing that antibodies to VEGF receptor can inhibit angiogenesis and thereby halt tumor progression. In accordance with the present invention, a vaccine composition comprising at least an immunogenic portion of the extracellular domain of VEGF receptor coupled to or administered with an antigenic protein or peptide such a KLH can be used as a

vaccine to provide the same therapeutic benefits using an active as opposed to a passive approach.

Still a further non-limiting example of a transmembrane protein which can be used in whole or in part in the method of the invention is the IL-2 receptor. The IL-2 receptor is expressed on most T-cells malignancies, and there is a data showing that antibodies to the IL-2 receptor can be used in the treatment of T-cell malignancies and autoimmune disorders. In the present invention, a composition is made comprising at least an immunogenic portion of the extracellular domain of the IL-2 receptor (e.g., P55 or P75), coupled to or administered with an antigenic carrier protein or peptide such as KLH. and used as a vaccine.

The vaccine compositions of invention can be used alone or in combination (concurrently or sequentially) with drugs or chemotherapy agents that provide therapeutic benefit for the condition being treated. In the case of NHL, suitable chemotherapy agents which can be used in combination with the CD20 based vaccine include alkylating agents, anthrocyclines, cis-platinum, fludarabine, corticosteroids and vinca alkaloids. These same chemotherapy agents which might be used in combination with other vaccine compositions for other forms of cancer.

EXAMPLE 1

20

25

44 amino acid fragments of the extracellular domains of humans and murine CD20 (amino acids 136-179, Seq. ID Nos. 1 and 2) were synthesized using a solid-phase FMOC peptide synthesizer and coupled to KLH using the methodology described in the Pierce Catalog Protocol. The peptide coupled to KLH was then prepared for injection by formulation with QS-21 adjuvant. Balb/c mice were injected according to one of the following protocols on days 1, 8, 15, 22 and 50 of the experiment:

- A. Murine CD20 fragment-KLH with QS-21 adjuvant
- B. Human CD20 fragment-KLH with QS-21 adjuvant
- C. KLH with QS-21 adjuvant
- D. QS-21 adjuvant

30

5

- E. P190 (irrelevant protein) coupled to KLH with QS-21 adjuvant
- F. B3A2 (irrelevant peptide) coupled to KLH with QS-21 adjuvant.

The animals were sacrificed on day 62 of the experiment.

Serum samples from the mice were diluted 1:200 and evaluated by BSA-blocked ELISA using goat-anti-mouse antibody conjugated to alkaline phosphatase for antibodies which bind to human CD20, mouse CD20 and KLH. As shown in Figs 1A and B, mice injected with human CD20 coupled to KLH (Fig. 1A) or mouse CD20 coupled to KLH (Fig. 1B) administration of xenogeneic antibody produced a significant polyclonal antibody response to both human and mouse CD20, while the response following administration of syngeneic antibody was principally limited to antibodies to the syngeneic form of CD20. Either xenogeneic or syngeneic peptide can therefore be used to generate an immune response.

To confirm the ability of the antibodies to bind to B cells, Raji cells (a form of human B-cell lymphoma that expresses CD20 on its surface) were blocked with human IgG, washed and then incubated for 30 minutes on ice with a 1:10 dilution of plasma from a mouse vaccinated with P-190-KLH control or huCD20-KLH. As a positive control, Raji cells were incubated with B1 antibody, or IgG2 as an isotypic negative control. After washing, the cells were incubated with goat-anti-mouse antibody, washed and fixed with 1% paraformaldehyde. Flow cytometry analysis was performed in a Becton-Dickinson FACScan. The results are shown in Figs 2A and 3B, wherein the shaded data set are the experimental data set and the outlined data set is the negative controls. As is apparent, there is a strong binding of mouse antibodies and Raji cells, comparable to that observed with B1 antibody.

25 <u>EXAMPLE 2</u>

To assess the number of B cells present in vaccinated mice, an evaluation was made of cells expressing CD19, a standard phenotypic marker for B cells. Spleens were harvested from the animals vaccinated in Example 1 and put into a single-cell suspension.

After counting the total number of cells, the cells were stained with FITC-labeled anti-mouse CD19 and the samples were analyzed by flow cytometry with a FACScan. 10,000 events

were collected. The percentage of CD19 positive cells minus the control gate was multiplied by the total number of cells to determine the number of CD19 positive cells in mice treated with the mouse and human CD20 peptide conjugates, and the P190 irrelevant peptide conjugate control.

5

As shown in Fig. 3, the absolute number of CD19 positive cells was significantly reduced in mice treated with either of the CD20 peptide conjugates. The level of CD19 positive cells is a reflection of the number of CD20 positive B cells, and the number of immature CD19⁺, CD20⁻ B cells in the samples. The absolute number of CD19⁺ B cells actually underestimates the therapeutic efficacy of the treatment for elimination of CD20⁺ B cells, however, since CD19 is expressed on B cell progenitor cells before expression of CD20.

EXAMPLE 3

Mice were injected five times over two months with one of four treatment protocols as follows:

human CD20 (44 aa fragment)-KLH plus QS1 human CD20 (44 aa fragment)-KLH human CD20 (44 aa fragment) plus QS21

KLH plus QS21

20

25

30

Blood was collected on week 9 for analysis by ELISA. Sera from the vaccinated mice were diluted 1:200 and incubated on BSA blocked plates coated with msCD20, huCD20, P190 or KLH. Secondary goat anti-mouse antibody conjugated to alkaline phosphatase was added, and the color change of p-nitrophenyl phosphate substrate was measured at 405 nm. The results are summarized in Figs. 7A-D. In most instances the peptide, carrier protein and adjuvant are all needed for optimal response, although some responses were detected using less than all of the components.

EXAMPLE 4

Mice were vaccinated according to the schedule of Example 3 using one of four treatment protocols: human CD20 (44 aa fragment)-KLH plus QS21 adjuvant, mouse

5

CD20 (44 amino acid fragment)-KLH plus QS21, P190 (irrelevant protein)-KLH +QS21 and KLH and QS21 alone. Mouse serum samples were evaluated by ELISA for the presence of antibodies reactive with msCD20, huCD20, P190 and KLH. The results are shown in Figs. 6A-D. Antibodies generated in mice after vaccination with human or mouse-derived CD20 fragments are specific for the peptides used, yet are capable of inducing immunity to the corresponding peptide from other species.

EXAMPLE 5

Mice were vaccinated five times over two months with huCD20 fragment-KLH conjugate with no adjuvant or in combination with one of three adjuvants: QS21, BCG or Alum. Serum samples from the vaccinated mice were tested by ELISA. The results are summarized in Figs. 8A-D. QS21 was found to be the most effective of those tested.

EXAMPLE 6

To confirm the observations of Example 2, nucleated spleen cells were recovered by centrifugation in a density gradient from mice vaccinated with a CD20-KLH conjugate (human or mouse) in the presence of QS21 adjuvant. 1×10^6 cells from each mouse were incubated with 2 μg of rat anti-mouse CD19 FITC-labeled antibody or with isotope-matched FITC labeled rat antibody. Cells were washed, fixed and analyzed with a Becton Dickinson FACScaliber cytometer. Figs 9A-C, D-F and G-I show the results for three exemplary mice of each vaccination group. The decrease in the peak reflecting levels of CD19 positive spleen cells in each of the mice is apparent.

۳	_
	به
,	0
r	বে

Score (T _{1/2} of Dissociation of Molecule Containing this Subsequence)	1600 48 21.2 1600 60 89.4 28.7
HLA Molecule	Kd Kd A_0201 Kd Kd A_0201 A_0201
Peptide Sequence	NFIRAHTPYI FIRAHTPYI FLKMcSLNFI HFLKmRRLEL IYDCePSNSS LIQTSKPYV ELIQtSKPYV
Species	human human mouse mouse mouse mouse

CLAIMS


	1.	A method for active vaccination against autologous cells expressing
transmembrane	e pro	teins comprising administering to a patient a vaccine composition
comprising at 1	least	an immunogenic portion of the extracellular domain of the
transmembrane	e pro	tein, or a xenogeneic homolog thereof, coupled to or administered with an
carrier protein	effe	ctive to break tolerance to the transmembrane protein and a
pharmaceutica	lly a	cceptable adjuvant.
	2.	The method of claim 1, wherein the transmembrane protein is selected
from the group	con	sisting of CD20, Her2-neu, VEGF receptor, epidermal growth factor
receptor, the C	D19	molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-
glycoprotein.		
	3.	The method of claim 1, wherein the transmembrane protein is CD20.
	4.	The method of claim 1, wherein the vaccine composition comprises a
peptide having	g the	sequence given by Seq. ID No 1 or 2.
	5.	The method claim 1, wherein the carrier protein is keyhole limpet
hemocyanin.		
	6.	The method of claim 5, wherein the transmembrane protein is selected
from the grou	рсо	nsisting of CD20, Her2-neu, VEGF receptor, epidermal growth factor
receptor, the (CD19	9 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-
glycoprotein.		


7. The method of claim 5, wherein the transmembrane protein is CD20.


- 8. The method of claim 7, wherein the vaccine composition comprises a peptide having the sequence given by Seq. ID No 1 or 2.
- 9. A method for active vaccination against B cells expressing CD20 comprising administering to a patient a vaccine composition comprising at least an immunogenic portion of the extracellular domain of CD20, or a xenogeneic homolog thereof, coupled to or administered with an carrier protein effective to break tolerance to the transmembrane protein and a pharmaceutically acceptable adjuvant.
- 10. The method claim 9, wherein the carrier protein is keyhole limpet hemocyanin.
- 11. The method of claim 9, wherein the vaccine composition comprises a peptide having the sequence given by Seq. ID No 1 or 2.
- 12. A method for treatment of B cell non-Hodgkin's lymphoma, comprising administering to a patient suffering from B cell non-Hodgkin's lymphoma a vaccine composition comprising at least an immunogenic portion of the extracellular domain of CD20, or a xenogeneic homolog thereof, coupled to or administered with an carrier protein effective to break tolerance to the transmembrane protein and a pharmaceutically acceptable adjuvant.
- 13. A vaccine composition comprising at least an immunogenic portion of the extracellular domain of the transmembrane protein, or a xenogeneic homolog thereof, coupled to or administered with an carrier protein effective to break tolerance to the transmembrane protein and a pharmaceutically acceptable adjuvant.
- 14. The composition of claim 13, wherein the transmembrane protein is selected from the group consisting of CD20, Her2-neu, VEGF receptor, epidermal growth

3	factor receptor	r, the C	D19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the
4	P-glycoprotein	n.	
1		15.	The composition of claim 13, wherein the transmembrane protein is
2	CD20.		
1		16.	The composition of claim 15, wherein the vaccine composition
2	comprises a p	eptide l	having the sequence given by Seq. ID No 1 or 2.
2		17.	The composition of claim 13, wherein the carrier protein is keyhole
<u>1</u> 2	limpet hemoc	yanin.	
Ample Mary Mary Mary Mary Mary Mary Mary Mary			
1		18.	The composition of claim 17, wherein the transmembrane protein is
<u></u> 2	selected from	the gro	oup consisting of CD20, Her2-neu, VEGF receptor, epidermal growth
3	factor recepto	r, the C	CD19 molecule, interleukin-2-receptor, interleukin-4-receptor, and the P-
3 4	glycoprotein.		
And the second s			
1		19.	The composition of claim 17, wherein the transmembrane protein is
2	CD20.		
1		20.	The composition of claim 19, wherein the vaccine composition

comprises a peptide having the sequence given by Seq. ID No 1 or 2.








TOTAL EVENTS: 10000

MARKER	LEFT, I	RIGHT	EVENTS	% TOTAL
ALL		9910	10000	100.00
M1	3	17	7525	75.25
M2	1.	3	1201	12.01
141 (' ' '	_		

FIG. 2A



TOTAL EVENTS: 10000

MARKER	LEFT, RIGHT	EVENTS_	% TOTAL
ALL	1, 9910	10000	100.00
M1	3, 12	7327	73.27
M2	1 4	1454	1 4.54

FIG. 2B

09/37/970






FIG. 3



SUBSTITUTE SHEET (RULE 26)







FIG. 6A

FIG. 6B






FIG. 6C

FIG. 6D




FIG. 7A




FIG. 7B




FIG. 7C




FIG. 7D




FIG. 8A




FIG. 8B




FIG. 8C






FIG. 8D





SEQUENCE LISTING

```
<110> Agus, David B.
        Scheinberg, David
        Zelenetz, Andrew D.
        Roberts, Wenay
  <120> Compositions and Methods for Active Vaccination
  <130> MSKP039US
  <140>
 <141>
 <150> 60/084,370
 <151> 1998-05-08
 <160> 6
 <170> PatentIn Ver. 2.0
<210> 1
 <211> 44
 <212> PRT
 <213> HUMAN
 <220>
 <223> human CD20 fragment
 <400> 1
 Lys Ile Sor His Phe Lou Lys Met Glu Ser Lou Asn Pho Ile Arg Ala
        5
                     10
His Thr Pro Tyr Ile Asn Ile Tyr Asn Cys Glu Pro Ala Asn Pro Ser
           20 25
                                     30
Glu Lys Asn Ser Pro Ser Thr Gln Tyr Cys Tyr Ser
        35
<210> 2
<211> 44
<212> PRT
<213> Murine
<220>
<223> murine CD20 fragment
```

```
<400> 2
   Thr Leu Ser His Phe Lou Lys Met Arg Arg Leu Glu Leu Ile Gln Thr
                                       10
   Ser Bys Pro Tyr val App The Tyr App Cys Glu Pro Ser Ash Ser Ser
                                   25
  Glu Lys Asn Ser Pro Ser Thr Gln Tyr Cys Asn Ser
           35
                              40
 <210> 3
  <211> 79
 <212> PRT
 <213> HUMAN
 <220>
 <223> exon VI of numan CD20 gene
 <400> 3
 Val Lys Gly Lys Met Ile Met Asn Ser Lou Ser Leu Phe Ala Ala Ile
                   5
                                      10
 Ser Gly Met Ile Leu Ser Ile Met Asp Ile Leu Asn Ile Lys Ile Ser
             20
                                  25
                                                      30
 His Pho Lou Lys His Glu Ser Leu Asn Phe Ile Arg Ala His Thr Pro
                              40
 Tyr Ile Asn Ile Tyr Asn Cys Glu Pro Ala Asn Pro Ser Glu Lys Asn
      50
                         55
 Ser Pro Ser Thr Glm Tyr Cys Tyr Ser Ile Gln Ser Leu Phe Leu
 65
                     70
<210> 4
<211> 237
<212> DNA
<213> HUMAN
<220>
<223> exon VI of human CD20 gene
<400> 4
gtcaaaggaa aaatgataat gaatteattg ageetetttg etgecattte tggaatgatt 60
ctttcaatca tggacatast taatattaaa atttcccatt ttttaaaaat ggagagtctg 120
aattttatta gageteasac accatatatt aacatataca actgtgaacc agetaateec 180
```

totgaganda actoccoato tacocantao tgttacagea tacaatotot gttottg	237
<210> 5 <211> 1255 <212> PRT <213> HUMAN	
<220> <223> numan Herî	
<400> 5 Met Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Leu Ala Leu Leu 1 5 10 15	
Pro Pro Cly Ala Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys 20 25 30	
Lou Arg Leu Pro Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His 35 40 45	
Leu Tyr Gln Gly Cys Gln Val Val Gln Gly Acn Leu Glu Leu Thr Tyr 50 55 60	
Leu Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val 65 70 75 80	
Gln Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu 85 90 95	
Gln Arg Lou Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr 100 105 110	
Ala Leu Ala Val Leu Asp Asn Gly Asp Pro Leu Asn Asa Thr Thr Pro 115 120 125	
Val Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser 130 135 140	
Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln 145 150 155 160	
Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn 135 170 175	
Asn Gln Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys 180 185 190	
His Pro Cys Ser Pro Met Cys Lys Gly Ser Arg Cys Trp Gly Glu Ser	

195		200	205	
Ser Glu Asp C 210	ys Gln Ser Leu 215	Thr Arg Thr V	7al Cys Ala Gl 220	y Gly Cy:
Ala Arg Cys L 225	ys Gly Pro Lcu 230		dys Cys His Gl 135	u Gln Cys 240
Ala Ala Gly C	ys Thr Gly Pro 245	Lys His Sor A 250	sp Cys Leu Ala	e Cys Lei 255
His Phe Asn Hi 26	is Ser Gly Ilc	Cys Glu Leu H. 265	is Cys Pro Ala 270	
Tnr Tyr Asn Th 275	er Asp Thr Phe	Glu Sor Met Pi 280	ro Asn Pro Glu 285	Gly Arg
Tyr Thr Pne Gl 290	y Ala Ser Cys V 295	Val inm Ala Cy	/s Pro Tyr Asn 300	Tyr Leu
Ser Thr Asp Va 305	l Gly Ser Cys 1 310	Chr Leu Val Cy 31		Asn Gln 320
Glu Val Thr Al	a Glu Asp Gly T 325	hr Gln Arg Cy 330	s Glu Lys Cys	Ser Lys 335
Pro Cys Ala Arg		ly Led Gly Me 345	t Glu His Leu 350	Arg Glu
Val Arg Ala Val 355		sn Ile Gln Gli 60	u Pho Ala Gly 365	Cys Lys
Lys Ile Phe Gly 370	Ser Leu Ala Pl 375	ne Leu Pro Glu	Ser Phe Asp	Gly Asp
Pro Ala Ser Asn 385	Thr Ala Pro Le	eu Glm Pro Glu 395		Val Phe 400
Glu Thr Leu Glu	Glu Ile Thr Gl 405	y Tyr Leu Tyr 410		Frp Pro 115
Asp Ser Leu Pro 420	Asp Leu Ser Va	l Pne Gln Asn 425	Leu Glm Val I 430	[le Arg
Gly Arg Ile Leu 435	His Asn Gly Ald		Thr Leu Gln G	ly Leu
Gly Ile Ser Trp	Leu Gly Leu Arg	g Ser Leu Arg	Glu Leu Gly S	er Gly

450 455 460
Leu Ala Leu Ile His His Asn Thr His Leu Cys Phe Val His Thr Val 465 470 475 480
Pro Trp Asp Gln Leu Pne Arg Asn Pro His Gln Ala Leu Lou His Thr 485 490 495
Ala Asn Arg Pro Glu Asp Glu Cys Val Gly Glu Gly Leu Ala Cys His 500 505 510
Glm Leu Cys Ala Arg Gly His Cys Trp Gly Pro Gly Pro Thr Glm Cys 515 520 525
Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys Val Glu Glu Cys 530 540
Arg Val Leu Glm Gly Leu Pro Arg Glu Tyr Val Asm Ala Arg His Cys 545 550 555 560
Lou Pro Cys His Pro Glu Cys Gln Pro Gln Asn Gly Ser Val Thr Cys 565 570 575
Phe Gly Pro Glu Ala Asp Gln Cys Val Ala Cys Ala His Tyr Lys Asp 580 585 590
Pro Pro Phe Cys Val Ala Arg Cys Pro Ser Gly Val Lys Pro Asp Leu 595 600 605
Ser Tyr Met Pro Ile Trp Lys Phe Pro Asp Glu Glu Gly Ala Cys Gln 610 615 620
Pro Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Asp Lys 625 630 635 640
Gly Cys Pro Ala Glu Gln Arg Ala Ser Pro Leu Thr Ser Ile Ile Ser 645 650 655
Ala Val Val Gly Ile Leu Leu Val Val Val Leu Gly Val Val Phe Gly 660 665 670
Ile Leu Ile Lys Arg Arg Gln Gln Lys Ile Arg Lys Tyr Thr Met Arg 675 680 685
Arg Leu Leu Gln Glu Thr Glu Leu Val Glu Pro Leu Thr Pro Ser Gly 690 . 695 700
Ala Met Pro Asn Gln Ala Gln Met Arg Ile Leu Lys Glu Thr Glu Leu

WO 99/57981 6 PCT/US99/10065

705	710	715	720
Arg Lys Val Lys Val	Lou Gly Ser Gly Ala		Tyr Lys
725	730		735
Gly Ile Trp Ile Pro 7	Asp Gly Glu Asn Val 745	Lys Ile Pro Val A	Ala Ile
Lys Val Leu Arg Glu <i>F</i>	Asn Thr Ser Pro Lys	Ala Asn Lys Glu I	le Leu
755	760	765	
Asp Glu Ala Tyr Val M	et Ala Gly Val Gly	Ser Pro Tyr Val S	er Arg
770	775	780	
Leu Leu Gly Ile Cys L		Gln Leu Val Thr G	ln Leu
785 7		795	800
Met Pro Tyr Gly Cys L	eu Leu Asp His Val <i>I</i>		ly Arg
805	810		15
Leu Gly Ser Glm Asp Lo	eu Leu Asn Trp Cys M	et Gln Ile Ala Ly	ys Gly
820	825	830	
Met Ser Tyr Leu Glu As	sp Val Arg Leu Val H	iis Arg Asp Leu Al	a Ala
835	840	845	
Arg Asn Val Leu Val Ly	rs Ser Pro Asn His Vo	al Lys Ile Thr As	p Phe
850	855	860	
Gly Leu Ala Arg Leu Le		hr Glu Tyr His Al.	a Asp
365 87		75	880
Gly Gly Lys Val Pro Ile	e Lys Trp Met Ala Lo	ou Glu Ser Ile Lei	
885	890	899	
Arg Arg Phe Thr His Glr	n Ser Asp Val Trp Se	er Tyr Gly Val Thr	: Val
900	905	910	
Trp Glu Lou Met Thr Phe	e Gly Ala Lys Pro Ty 920	r Asp Gly Ile Pro	Ala
Arg Glu Ile Pro Asp Leu	Leu Glu Lys Gly Glu	u Arg Leu Pro Gln	Pro
930	935	940	
Pro Ile Cys Thr Ile Asp 945 950	Val Tyr Met Ile Met		Met 960
Ile Asp Ser Glu Cys Arg	Pro Arg Phe Arg Glu	ı Leu Val Ser Glu	Phe

965		976	975
Ser Arg Met Ala Arg 980	Asp Pro Gln Arg 985	Phc Val Val Ile	Gln Asn Glu 990
Asp Leu Gly Pro Ala 995	Ser Pro Leu Asp	Sor Thr Phe Tyr 1005	Arg Scr Leu
Leu Glu Asp Asp Asp 1010	Met Gly Asp Leu 1 1015	/al Asp Ala Glu 1020	Glu Tyr Leu
Val Pro Gln Gln Gly 1025 1	Phe Phe Cys Pro A 030	asp Pro Ala Pro 1035	Gly Ala Gly 1040
Gly Not Val His His A		er Sor Thr Arg 50	Ser Gly Gly 1055
Gly Asp Lou Thr Leu (1060	Gly Leu Glu Pro S 1065		Ala Pro Arg 070
Ser Pro Leu Ala Pro S 1075	Ser Glu Cly Ala G 1080	ly Ser Asp Val 1085	Phe Asp Gly
Asp Leu Gly Met Gly A 1090	la Ala Lys Gly Lo 1095	eu Gln Ser Leu 1 1100	Pro Thr His
Asp Pro Ser Pro Leu G		u Asp Pro Thr (/al Pro Leu 1120
Pro Ser Glu Thr Asp G	ly Tyr Val Aia Pr 113		Ser Pro Gln 1135
Pro Glu Tyr Val Asn Gl	ln Pro Asp Val Ar 1145	g Pro Gln Pro P 11	
Arg Glu Gly Pro Leu Pr 1155	o Ala Ala Arg Pro 1160	o Ala Gly Ala T 1165	hr Leu Glu
Arg Pro Lys Thr Leu Sc	r Pro Gly Lys Asr 1175	n Gly Val Val Ly 1180	ys Asp Val
Phe Ala Phe Gly Gly Al 1185 119		Glu Tyr Lou Th	er Pro Gln 1200
Gly Gly Ala Ala Pro Gli 1205	n Pro His Pro Pro 1210		r Pro Ala 1215
Phe Asp Asm Leu Tyr Tyr	Trp Asp Gln Asp	Pro Pro Glu Ar	g Gly Ala

WO 99/57981 PCT/US99/10065

8

1220 1225 1230

Pro Pro Ser Thr Phe Lys Gly Thr Pro Thr Ala Glu Asn Pro Glu Tyr 1235 1240 1245

Leu Gly Leu As; Val Pro Val 1250 1255

<210> 6

<211> 1210

<212> PRT

<213> HUMAN

<220>

<223> numan EGFR

<400> 6

Mot Arg Pro Scr Gly Thr Ala Gly Ala Ala Leu Leu Ala Leu Leu Ala 1 5 10 15

Ala Leu Cys Pro Ala Ser Arg Ala Leu Glu Glu Lys Lys Val Cys Gln
20 25 30

Gly Thr Ser Asn Lys Leu Thr Gln Leu Gly Thr Phe Glu Asp His Phe 35 40 45

Leu Ser Leu Gln Arg Met Phe Asn Asn Cys Glu Val Val Leu Gly Asn
50 60

Leu Glu Ile Thr Tyr Val Gln Arg Asn Tyr Asp Leu Ser Phe Leu Lys
65 70 75 80

Thr IIc Gln Glu Val Ala Gly Tyr Val Leu IIc Ala Leu Asn Thr Val
85 90 95

Glu Arg Ile Pro Leu Glu Asn Leu Gln Ile Ile Arg Gly Asn Met Tyr 100 105 110

Tyr Glu Asn Ser Tyr Ala Leu Ala Val Leu Ser Aon Tyr Asp Ala Asn 115 120 125

Lys Thr Gly Leu Lys Glu Leu Pro Met Arg Asn Leu Gln Glu Ile Leu 130 135 140

His Gly Ala Val Arg Phe Ser Asn Asn Pro Ala Leu Cys Asn Val Glu 145 150 155 160

Ser Ile	e Gln Tr	p Arg As; 165	o Ile V		Ser Asp P. 170	he Leu Se	er Asn Met 175
Ser Met	Asp Pho		n His D	eu Gly S 185	Ger Cys G	ln Lys Cy 19	s Asp Pro 0
Ser Cys	Pro Asi 195	n Gly Ser	Cys T: 2(la Gly Gl	u Gla As 205	n Cys Gln
Lys Leu 210		: Ile Ile	Cys Al 215	la Gln G	ln Cys Se 22		g Cys Arg
Gly Lys 225	Ser Pro	Ser Asp 230	Cys Cy	s His A	sn Gln Cy 235	s Ala Ala	Gly Cys 240
Thr Gly	Pro Arg	Glu Ser 245	Asp Cy	s Lou Va 25	al Cys Ar	g Lys Pho	e Arg Asp 255
Glu Ala	Thr Cys 260	Lys Asp	Thr Cy	s Pro Pr 265	ro Leu Mel	Leu Tyr 270	
	Tyr Gln 275	Met Asp	Val Asr 280		u Gly Lys	: Tyr Ser 285	Phe Gly
Ala Thr 290	Cys Val		Cys Pro 295	Arg As	n Tyr Val 300		Asp His
Gly Ser	Cys Val	Arg Ala 310	Cys Gly	· Ala As _j	p Ser Tyr 315	Glu Met	Glu Glu 320
Asp Gly '		Lys Cys : 325	pa pa	Cys Glu	ı Gly Pro)	Cys Arg	Lys Val 335
Cys Asn (Gly Ile	Gly Ile (Gly Glu	Phe Lys	s Asp Ser	Leu Ser 350	Ile Asn
Ala Thr A	Asn Ile 1 155	Lys His A	he Lys 360	Asn Cys	Thr Ser	Ile Ser 365	Gly Asp
Leu His I 370	le Leu 3		la Phe 75	Arg Gly	Asp Ser 380	Phe Thr	His Thr
Pro Pro L 385	eu Asp F	ro Gln G 390	lu Leu	Asp Ile	Leu Lyc 395	Thr Val I	Lys Glu 400
Ile Thr G	-	cu Leu I 05	le Gln	Ala Trp 410	Pro Glu .		Chr Asp 115

I	eu	Hli	ű A		Phe 120	Glu	Asr	. Le	u G		11e 425	īle	s yı	rg Gl	ly i	Arg	Thr 430		s	Gln
13	LS	Gly		ln I 35	he.	Ser	Leu	λla		al ' 40	Va!	Ser	î LiC	eu As		Ile M5	Thr	Se	r	Leu
G	_	Let 450		rg S	er.	Leu	Lys	Glu 455		le S	Ser	Asp	Gl	у Лs 46		al	Ile	īl	e S	Ser
	ly . 55	Asn	. Ly	s A	sn I	Leu	Cys 470	Tyr	· A3	a P	sn	Thr	Il 47	e As 5	n T	rp	Lys	Lys		eu 80
Pì	ie (Gly	Th	r So		31 <i>y</i> 85	Gln	Lys	Th	r L		Ile 490	Ile	e Se:	r A	sn .	Arg	Gl _y 495		lu
ХS	n S	Ser	Су	s Ly 50		la '	Thr	Gly	Gl		al (05	Сув	Нів	s Alá	ı Lo		Cys 510	Ser	P.	ro
Gl	u G	ly	Су: 515		p G	ly 1	Pro	Glu	Pro		rg A	/sp	Суз	Val	. S∈ 52		Cys	Arg	A	sn
Va		er 30	Arq	g Gl	y Ai	rg (Cys 535	Val	L As	sp L	ys	Cys	Lys 540		u L	eu (Glu	G3	.У
Gl: 545		ro	Arg	r Gl	u Př		/al (50	Glu	Asn	ı Se	r G		Cys 555	Ile	G1:	n C	ys 1	lis	Pr 56	
Glu	ı C)	/s :	Leu	Pro	5 Cl		la M	iet .	Asn	I1		hr (Сув	Thr	Gly	y A	-	75	Pr	o
Asp	As	sn (Суз	Ilc 580		n Cj	ys A	la i	His	Ту: 58!		le 7	(as	Gly	Pro		15 C	λa	Va:	l
Lys	Th		Cys 595	Pro	Al.	a Gi	ly V		Met 500	Gly	/ G]	Lu A	sn	Asn	Thr		eu V	al	Trį	>
Lys	Ту 61		.la	Asp	Ala	a Gl		is V 15	'al	Cys	H1	s L		Cys 620	Hıs	Pr	o A:	sn (Cys	;
Thr 625	Ту:	r G	ly	Суъ	Tnr	Gl 63	_	co G	ly	Leu	Gl		ly (35	Cys .	Pro	Th	r As		51y 540	
Pro	Lys	зI	le	Pro	Ser 545	I1	e Al	a Tì	hr	Gly	Me:		al C	Gly A	Ala	Lei	ս L∈ 65		eu	
Leu	Let	. Vá		Val	Ala	Lei	u Gl	y I		Sly	Let	a Ph	ne M	let /	ırg	Arg		g H	ıs	

Ile Val Arg Lys Arg Thr Leu Arg Arg Leu Leu Gln Glu Arg Glu Leu 675 680 685
Val Glu Pro Leu Thr Pro Ser Gly Glu Ala Pro Asn Gln Ala Leu Leu 690 695 700
Arg Ilc Leu Lys Glu Thr Glu Phe Lys Lys Ile Lys Val Leu Gly Ser 705 710 715 720
Gly Ala Phe Gly Thr Val Tyr Lys Gly Leu Trp Ile Pro Glu Gly Glu T25 730 735
Lys Val Lys Ile Pro Val Ala Tle Lys Glu Leu Arg Glu Ala Thr Ser 740 745 750
Pro Lys Ala Ash Lys Glu Ile Lou Asp Glu Ala Tyr Val Met Ala Ser 755 760 765
Val Asp Asn Prc His Val Cys Arg Leu Leu Gly Ile Cys Leu Thr Ser 770 775 780
Thr Val Gln Leu Ile Thr Gln Leu Met Pro Phe Gly Cys Leu Leu Asp 785 790 795 800
Tyr Val Arg Glu His Lys Asp Asn Ile Gly Ser Gln Tyr Leu Leu Asn 805 810 815
Trp Cys Val Glm Ile Ala Lys Gly Met Asn Tyr Leu Glu Asp Arg Arg 826 825 830
Leu Val His Arg Asp Leu Ala Ala Arg Ash Val Leu Val Lys Thr Pro 835 840 845
Gln His Val Lys Ile Thr Asp Phe Gly Leu Ala Lys Leu Leu Gly Ala 850 855 860
Glu Glu Lys Glu Tyr His Ala Glu Gly Gly Lys Val Pro Ile Lys Trp 865 870 875 880
Met Ala Leu Gl. Ser Ile Leu His Arg Ile Tyr Thr His Gln Ser Asp 385 890 895
Val Trp Ser Tyr Gly Val Thr Val Trp Glu Leu Met Thr Phe Gly Ser 900 905 910
Lys Pro Tyr Asp Gly Ile Pro Ala Ser Glu Ile Ser Ser Ile Leu Glu 915 920 925

Lys Gly Glu Arg Leu Pro Gln Pro Pro Ile Cys Thr Ile Asp Val Tyr 920 935 940

Met Ile Met Val Lys Cys Trp Met Ile Asp Ala Asp Sar Arg Pro Lys 945 950 955 960

Phe Arg Glu Leu Ile Ile Glu Phe Ser Lys Met Ala Arg Asp Pro Gln 965 970 975

Arg Tyr Lou Val Ile Gln Gly Asp Glu Arg Met His Leu Pro Ser Pro 980 985 990

Thr Asp Ser Ash Phe Tyr Arg Ala Leu Met Asp Glu Glu Asp Met Asp 995 1000 1005

Asp Val Val Asp Ala Asp Glu Tyr Leu Ile Pro Gln Gln Gly Phe Phe 1010 1015 1020

Ser Ser Pro Ser Thr Ser Arg Thr Pro Leu Leu Ser Ser Leu Ser Ala 1025 1030 1035 1040

Thr Ser Asn Asn Ser Thr Val Ala Cys Ile Asp Arg Asn Gly Leu Gln 1045 1050 1055

Ser Cys Pro Ile Lys Glu Asp Ser Phe Leu Gln Arg Tyr Ser Ser Asp 1060 1065 1070

Pro Thr Gly Ala Leu Thr Glu Asp Ser Ile Asp Asp Thr Phe Leu Pro 1075 1080 1085

Val Pro Glu Tyr Ile Asn Gln Ser Val Pro Lys Arg Pro Ala Gly Ser 1090 1095 1100

Val Gln Asn Pro Val Tyr His Asn Gln Pro Leu Asn Pro Ala Pro Ser 1105 1110 1115 1120

Arg Asp Pro His Tyr Gln Asp Pro His Ser Thr Ala Val Gly Asn Pro 1125 1130 1135

Glu Tyr Leu Asn Thr Val Gln Pro Thr Cys Val Asn Ser Thr Phe Asp 1140 1145 1150

Ser Pro Ala His Trp Ala Gln Lys Gly Scr His Gln Ile Ser Leu Asp 1155 1160 1165

Ash Pro Asp Tyr Gln Gln Asp Phe Phe Pro Lys Glu Ala Lys Pro Ash 1170 1175 1180

Gly Ile Phe Lys Gly Ser Thr Ala Glu Asn Ala Glu Tyr Leu Arg Val 1185 1190 1195 1200

Ala Pro Gln Ser Ser Glu Phe Ile Gly Ala 1205 1210

OPPEDAHL & LARSON

FILE NO. MSKP039 INVENTOR . Agus, et al

Claim for Priority

I hereby claim foreign priority benefits under 35 U.S.C. § 119 (a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below any foreign applications for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)	PRIORITY CLAIMED	CERTIFIED COPY ATTACHED
·		· <u>-</u>		YES[]NO[]	YES[]NO[]
FOREIGN APPLICA	ATION(S), IF ANY, FILED MO	ORE THAN 12 MONTH	IS (6 MONTHS FOR I	DESIGN) PRIOR T	O SAID
COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)		

I hereby claim the benefit under 35 U.S.C § 119(e) of any United States provisional application(s) listed below.

60/084,870	08 May 1998	
(application number)	(filing date)	
(application number)	(filing date)	

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF SOLE OR FIRST INVENTOR	LAST NAME AGUS	FIRST NAME David	MIDDLE NAME B.
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE Brooklyn Beverly Hills PM	STATE OR COUNTRY OF RESIDENCE	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRESS PICTOPONT Street		CITY Brooklyn DA Berey Hills	STATE/COUNTRY ZIP CODE NY 10021 PA CA 90210
DATE ///)/		SIGNATURE	

[X] Signature for additional joint inventor attached. Number of Pages 1. [] Signature by Administrator(trix) or legal representative for deceased or

incapacitated inventor. Number of Pages ___.

[] Signature for inventor who refuses to sign or cannot be reached by person authorized under 37 CFR § 1.47. Number of Pages ___.

Ь∀@E 03\0⊄

FILE NO. <u>MSKP039</u> INVENTOR . Agus, et al

NAME OF SECOND INVENTOR	LAST NAME SCHEINBERG	FIRST NAME DAVID	MIDDLE NAME	
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE NEW YORK	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US	
POST OFFICE ADDRE 325 Central Park \		CITY New York	STATE/COUNTRY ZIP CODE NY 10025	
DATE	Co	SIGNATURE)	
NAME OF THIRD INVENTOR	LAST NAME ROBERTS	FIRST NAME WENDY	MIDDLE NAME	
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE NEW YORK N	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US	
POST OFFICE ADDRE 1233 York Avenue	ess 303East71St	CITY New York	STATE/COUNTRY ZIP CODE NY 10021	
DATE / /DE)	SIGNATURE WHILLIAM PHI	5	
NAME OF FOURTH INVENTOR	LAST NAME ZELENETZ	FIRST NAME ANDREW	MIDDLE NAME D.	
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE LARCHMONT	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US	
POST OFFICE ADDRE 31 Mohegan Road		CITY Larchmont	STATE/COUNTRY ZIP CODE NY 10538	
DATE 11/6/2005	5	SIGNATURE Sun telem		

FILE NO. MSKP039 INVENTOR . Agus, et al

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My citizenship, residence and post office address are as listed below next to my name.

l believ which a	e I am the origina a patent is sought	l, first and [] sole/[i on the invention er	x] joint inventor of the subject m ntitled: <u>Compositions and Metho</u>	atter which is claimed and design to the design at the des	d for
the spe	ecification of which	1			
(a)[]	is attached herel				
(b)[]	was filed on on	as	s Application Serial No.	and was ame	ended
(c) [X]	was described a and amended or	nd claimed in Inter	national Application No. <u>PCT/U</u>	<u>S99/10065</u> filed on <u>May 7</u>	<u>, 1999</u>
includi	ing the claims, as a ation which is mat	e reviewed and und amended by any a erial to the patenta	edgment of Duty of Disclosure derstood the content of the above mendment referred to above. I ability of the subject matter claim Regulations § 1.56(a).	e identified specification, acknowledge the duty to	disclose
365(c) insofar States	of any PCT interr r as the subject ma for PCT internation wledge the duty to en the filing date of	ational application atter of each of the nal application in the disclose material	35 U.S.C. § 120 nited States Code, § 120 of any designating the United States of claims of this application is not the manner provided by the first information as defined in 37 CF ion and the national or PCT into	disclosed in the prior Uniperagraph of 35 U.S.C. §	ited 112, l available
(Applica	ation Serial No.)	(Filing Date)	(Status)(patented,pending,abando	ned) (Patent No. if a	oplicable)
(Applica	ation Serial No.)	(Filing Date)	(Status)(patented,pending,abando	ned) (Patent No. if a	pplicable)
			Power of Attorney		
Nancy	/ J. Parsons, PTO	Reg. No. 40,364 o Center 2 nd Floor 2	. NO. 32,746, Marina T. Larson of the firm of OPPEDAHL & LAR 256 Dillon Ridge Rd., Dillon, CO os in the Patent and Trademark (80435 as attorneys to pr	osecute
SEN	D CORRESPONDENCE O21121	a.d.	DIRECT TELEPHONE O OPPEDAHL & LARSON (970) 468-6600		

PATENT TRADEHMAK OFFICE

FILE NO. MSKP039 INVENTOR . Agus, et al

Claim for Priority

I hereby claim foreign priority benefits under 35 U.S.C. § 119 (a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below any foreign applications for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

EARLIEST FOREIGN APPLICATION	APPLICATION(S), FILED	WITHIN TWELVE MO	NIHS (6 MONIHS F	UR DESIGN) PRIC	OR TO SAID
COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)	PRIORITY CLAIMED	CERTIFIED COPY ATTACHED
				YES[]NO[]	YES[]NO[]
FOREIGN APPLICAT	TON(S), IF ANY, FILED MO	ORE THAN 12 MONTH	IS (6 MONTHS FOR I	DESIGN) PRIOR T	O SAID
COUNTRY	APPLICATION NO.	DATE OF FILING (day/month/year)	DATE OF ISSUE (day/month/year)		
		Provisional App	olication		
hereby claim the lelow.	benefit under 35 U.S.	C § 119(e) of any	United States pro	visional applica	ation(s) listed
60/084,870		08 Ma	ay 1998		
(application number	er)	(filing d	ate)		
(application numbe	er)	(filing d	ate)		

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

NAME OF SOLE LAST NAME OR FIRST AGUS		FIRST NAME David	MIDDLE NAME B.
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE Brooklyn	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US
POST OFFICE ADDRE 9 Pierrepont Stree		CITY Brooklyn	STATE/COUNTRY ZIP CODE NY 10021
DATE		SIGNATURE	

- [X] Signature for additional joint inventor attached. Number of Pages 1
- [] Signature by Administrator(trix) or legal representative for deceased or incapacitated inventor. Number of Pages ___.
- [] Signature for inventor who refuses to sign or cannot be reached by person authorized under 37 CFR § 1.47. Number of Pages ___.

OPPEDAHL & LARSON

FILE NO. <u>MSKP039</u> INVENTOR . Agus, et al

NAME OF SECOND INVENTOR	LAST NAME SCHEINBERG	FIRST NAME DAVID	MIDDLE NAME		
RESIDENCE & CITIZENSHIP	CITY OF RESIDENCE NEW YORK	STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US		
POST OFFICE ADDRE		CITY New York STATE/COUNTRY ZIP CODE NY 10025			
DATE		SIGNATURE			
NAME OF THIRD INVENTOR	LAST NAME ROBERTS	FIRST NAME WENDY	MIDDLE NAME		
RESIDENCE & CITY OF RESIDENCE NEW YORK		STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US		
POST OFFICE ADDRE		CITY New York	STATE/COUNTRY ZIP CODE NY 10021		
DATE		SIGNATURE			
NAME OF FOURTH INVENTOR	LAST.NAME ZELENETZ	FIRST NAME ANDREW	MIDDLE NAME D.		
RESIDENCE & CITY OF RESIDENCE LARCHMONT		STATE OR COUNTRY OF RESIDENCE NY	COUNTRY OF CITIZENSHIP US		
POST OFFICE ADDRESS 31 Mohegan Road		CITY Larchmont STATE/COUNTRY ZIP CODE NY 10538			
DATE		SIGNATURE			

OPPEDAHL & LARSON

FILE NO. MSKP039 INVENTOR . Agus, et al

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My citizenship, residence and post office address are as listed below next to my name.

I believe I am the original, first and [] sole/[x] joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled Compositions and Methods for Active Vaccination

WHICH	a patent is sought o	n the invention e	ntitied:Compositions and Method	ds for Active Vaccination
the spe	ecification of which		•	
(a)[]	is attached hereto	·.		
(b) [¹]	was filed on	as	s Application Serial No.	and was amended
(c) [X]	was described and claimed in International Application No. <u>PCT/US99/10065</u> filed on <u>May 7, 1999</u> and amended on			
includir informa	ng the claims, as ar ation which is mate	reviewed and uncomended by any a rial to the patenta	edgment of Duty of Disclosure derstood the content of the above mendment referred to above. I ability of the subject matter claim Regulations § 1.56(a).	e identified specification, acknowledge the duty to disclose
365(c) insofar States acknow	of any PCT interna as the subject mat or PCT international vedge the duty to den the filing date of	tional application ter of each of the al application in the tisclose material	35 U.S.C. § 120 nited States Code, § 120 of any designating the United States o claims of this application is not one manner provided by the first p information as defined in 37 CFF ion and the national or PCT inter	of America, listed below and, disclosed in the prior United paragraph of 35 U.S.C. § 112, I R § 1.56 which became available
(Applicat	tion Serial No.)	(Filing Date)	(Status)(patented,pending,abandon	ned) (Patent No. if applicable)
(Applicat	tion Serial No.}	(Filing Date)	(Status)(patented,pending,abandon	ned) (Patent No. if applicable)
			Power of Attorney	
Nancy Box 50	J. Parsons, PTO R 68, Alpine Bank Ce	eg. No. 40,364 o enter, 2 ^{na} Floo r , 2	NO. 32,746, Marina T. Larson, f the firm of OPPEDAHL & LARS 56 Dillon Ridge Rd., Dillon, CO & in the Patent and Trademark O	SON LLP, having office at P.O. 80435 as attorneys to prosecute
SEND	CORRESPONDENCE	TO:	DIRECT TELEPHONE CA OPPEDAHL & LARSON L (970) 468-6600	

PATENT TRACEWORK OFFICE